

# A REMARK ON THE CHEBOTAREV THEOREM ABOUT ROOTS OF UNITY.

F. PAKOVICH

**ABSTRACT.** Let  $\Omega$  be a matrix with entries  $a_{i,j} = \omega^{ij}$ ,  $1 \leq i, j \leq n$ , where  $\omega = e^{2\pi\sqrt{-1}/n}$ ,  $n \in \mathbb{N}$ . The Chebotarev theorem states that if  $n$  is a prime then any minor of  $\Omega$  is non-zero. In this note we provide an analogue of this statement for composite  $n$ .

Let  $\Omega$  be a matrix with entries  $a_{i,j} = \omega^{ij}$ ,  $1 \leq i, j \leq n$ , where  $\omega = e^{2\pi\sqrt{-1}/n}$ ,  $n \in \mathbb{N}$ . The Chebotarev theorem states that if  $n$  is a prime then any minor of  $\Omega$  is non-zero. Chebotarev's proof of this theorem and the references to other proofs can be found in [2]. Yet another proofs can be found in recent papers [1], [3].

For a complex polynomial  $P(z)$  denote by  $w(P)$  the number of non-zero coefficients of  $P(z)$ . It is easy to see that the Chebotarev theorem is equivalent to the following statement: if a non-zero polynomial  $P(z)$ ,  $\deg P(z) \leq n-1$ , has  $k$  different roots which are  $n$ -roots of unity then  $w(P) > k$  whenever  $n$  is a prime.

A natural question is: how small can be  $w(P)$  if  $n$  is a composite number? The example of the polynomial

$$D_{n,r,l}(z) = z^l(1 + z^r + z^{2r} + \dots + z^{(\frac{n}{r}-1)r}),$$

where  $r|n$ ,  $0 \leq l \leq r-1$ , shows that  $w(P)$  could be as small as  $n/(n-k)$ . In this note we show that actually it is the "worst" possible case.

**Theorem.** *Let  $n$  be a composite number and  $P(z)$  be a non-zero complex polynomial,  $\deg P(z) \leq n-1$ . Suppose that  $P(z)$  has exactly  $k$  different roots which are  $n$ -roots of unity. Then the inequality*

$$w(P) \geq \frac{n}{n-k} \quad (*)$$

*holds. Furthermore, the equality attains if and only if  $P(z)$  up to a multiplication by a complex number coincides with  $D_{n,r,l}(\omega^j z)$  for some  $j$ ,  $0 \leq j \leq n-1$ , and  $r, l$  as above.*

*Proof.* Let  $P(z) = p_0 + p_1 z + \dots + p_{n-1} z^{n-1}$  and let

$$C = \begin{pmatrix} p_0 & p_1 & \dots & p_{n-1} \\ p_{n-1} & p_0 & \dots & p_{n-2} \\ \dots & \dots & \dots & \dots \\ p_1 & p_2 & \dots & p_0 \end{pmatrix}$$

be the circulant matrix generated by the coefficients of  $P(z)$ . We will denote the row vectors of  $C$  by  $\vec{t}_j$ ,  $0 \leq j \leq n-1$ . Set  $r = \text{rk } C$ . The key observation is that

---

1991 *Mathematics Subject Classification.* 11T22, 11C08.

*Key words and phrases.* Chebotarev theorem, roots of unity, polynomials.

Research supported by the ISF, Grant No. 979/05.

the number  $k$  is equal to the number  $n - r$ . To establish it notice that eigenvectors of  $C$  are  $\vec{f}_i = ((\omega^i)^0, (\omega^i)^1, \dots, (\omega^i)^{(n-1)})$ ,  $0 \leq i \leq n - 1$ , and the corresponding eigenvalues are  $P(\omega^i)$ ,  $0 \leq i \leq n - 1$ . Furthermore, the vectors  $\vec{f}_i$ ,  $0 \leq i \leq n - 1$ , form a basis of  $\mathbb{C}^n$ . The matrix  $C$  is diagonal with respect to this basis and therefore  $k = n - r$ .

It follows that in order to prove inequality (\*) it is enough to establish the inequality

$$w(P)r \geq n. \quad (**)$$

This inequality essentially is a particular case of Theorem C in [1] and can be established as follows ([1]). Let  $V$  be a vector space generated by the vectors  $\vec{t}_j$ ,  $0 \leq j \leq n - 1$ , and  $R$  be a subset of  $\{\vec{t}_0, \vec{t}_1, \dots, \vec{t}_{n-1}\}$  consisting of  $r$  vectors which generate  $V$ . Clearly, for any  $i$ ,  $1 \leq i \leq n$ , there exists a vector  $v \in V$  for which its  $i$ -th coordinate is distinct from zero. Since each vector from  $R$  has exactly  $w(P)$  non zero coordinates it follows that (\*\*) holds.

For a vector  $\vec{v} \in \mathbb{C}^n$  denote by  $\text{supp}\{\vec{v}\}$  the set consisting of numbers  $i$ ,  $1 \leq i \leq n$ , for which  $i$ -th coordinate of  $\vec{v}$  is non-zero. Observe now that the equality in (\*\*) attains only if for any two vectors  $v_1, v_2 \in R$  we have  $\text{supp}\{v_1\} \cap \text{supp}\{v_2\} = \emptyset$ . This implies easily that  $\text{supp}\{\vec{t}_0\}$  consists of numbers congruent by modulo  $r$  to a number  $l$ ,  $0 \leq l \leq r - 1$ , and therefore  $P(z) = z^l Q(z^r)$  for some polynomial  $Q(z) = q_0 + q_1 z + \dots + q_{n/r-1} z^{n/r-1}$  and number  $l$ ,  $0 \leq l \leq r - 1$ .

Furthermore, since the vectors  $t_0, t_r, t_{2r}, \dots, t_{n/r-1}$  have equal supports the equality in (\*\*) implies that any two of them are proportional. Therefore, the rank of the circulant matrix  $W$  generated by the coefficients of  $Q(z)$  equals 1. This implies that the vector  $\vec{q} = \{q_0, q_1, \dots, q_{n/r-1}\}$  is orthogonal to  $n/r - 1$  vectors from the collection  $\vec{g}_j = ((\nu^j)^0, (\nu^j)^1, \dots, (\nu^j)^{(n/r)-1})$ ,  $0 \leq j \leq n/r - 1$ , where  $\nu = \omega^r$ . Since  $\vec{g}_j$ ,  $0 \leq j \leq n/r - 1$ , are linearly independent this implies that there exists  $\alpha \in \mathbb{C}$  such that  $\vec{q} = \alpha \vec{g}_j$  for some  $0 \leq j \leq n/r - 1$ .

#### REFERENCES

1. D. Goldstein, R. Guralnick, I. Isaacs, *Inequalities for finite group permutation modules*, Trans. Am. Math. Soc. 357, No.10, 4017-4042 (2005)
2. P. Stevenhagen, H. Lenstra, *Chebotarev and his density theorem*, Math. Intell. 18, No.2, 26-37 (1996)
3. T. Tao, *An uncertainty principle for cyclic groups of prime order*, Math. Res. Lett. 12, No.1, 121-127 (2005).

DEPARTMENT OF MATHEMATICS, BEN GURION UNIVERSITY, P.O.B. 653, BEER-SHEVA 84105, ISRAEL

E-mail address: pakovich@math.bgu.ac.il